



The Open Respiratory Medicine Journal

Content list available at: <https://openrespiratorymedicinejournal.com>

Supplementary Material



Current Utility of Sequential Organ Failure Assessment Score: A Literature Review and Future Directions

Rahul Kashyap¹ , Khalid M. Sherani^{2,3} , Taru Dutt⁴ , Karthik Gnanapandithan⁵ , Malvika Sagar⁶ ,
Saraschandra Vallabhajosyula⁷ , Abhay P. Vakil^{6,7}  and Salim Surani^{3,8,*} 

¹Department of Anesthesiology and Perioperative Medicine, Mayo Clinic, Rochester, MN 55905, USA

²Department of Internal Medicine, Jamaica Hospital Medical Center, Jamaica, NY 11418, USA

³Corpus Christi Medical Center, Corpus Christi, TX 78411, USA

⁴Department of Neurology, Mayo Clinic College of Medicine, Mayo Clinic, Rochester MN, USA and Hennepin County Medical Center, Minneapolis, MN 55905, USA

⁵Department of Internal Medicine, Yale-New Haven Hospital and Yale University School of Medicine, New Haven, CT 06510, USA

⁶Department of Pediatrics, McLane Children's Hospital, Baylor Scott and White Health, Temple, TX 76502, USA

⁷Critical Care Medicine, Mayo Clinic College of Medicine, Mayo Clinic, Rochester, MN 55905, USA

⁸Texas A&M University System Health Science Center, Bryan, TX 77807, USA

Article History

Received: October 2, 2020

Revised: December 13, 2020

Accepted: January 13, 2021

Table S1. Sequential Organ Failure Assessment score from original article [8].

	SOFA Score				
	0	1	2	3	4
Respiration	-	-	-	-	-
PaO ₂ /FiO ₂ (torr)	>400	≤400	≤300	≤200	≤100
	-	-	-	With respiratory support	With respiratory support
Coagulation	-	-	-	-	-
Platelets (x10 ³ /mm ³)	>150	≤150	≤100	≤50	≤20
Liver	-	-	-	-	-
Bilirubin (mg/dL)	<1.2	1.2-1.9	2.0-5.9	6.0-11.9	>12.0
(μmol/L)	<20	20-32	33-101	102-204	>204
Cardiovascular	-	-	-	-	-
Hypotension	No	MAP < 70mm Hg	Dopamine ≤6 or dobutamine (any dose)*	Dopamine > 6 or epi ≤ 0.1 or norepi ≤ 0.1*	Dopamine > 15 or epi > 0.1 or norepi > 0.1*
Central Nervous System	-	-	-	-	-
Glasgow Coma Score	15	13-14	10-12	6-9	<6
Renal	-	-	-	-	-
Creatinine (mg/dL)	<1.2	1.2-1.9	2.0-3.4	3.5-4.9	>5.0
(μmol/L)	<110	110-170	171-299	300-440	>440
or urine output	-	-	-	or <500 mL/day	or <200 mL/day

epi, epinephrine; norepi, norepinephrine.

*Adrenergic agents administered for at least 1 hr (doses are given in μg/kg/min)

To convert torr to kPa, multiply the value by 0.1333.

Table adapted from: Vincent JL, Moreno R, Takala J, Willatts S, De Mendonca A, Bruining H, Reinhart CK, Suter PM, Thijs LG: The SOFA (Sepsis-related Organ Failure Assessment) score to describe organ dysfunction/failure. On behalf of the Working Group on Sepsis-Related Problems of the European Society of Intensive Care Medicine. *Intensive Care Med* 1996, 22(7):707-710.

© 2021 Kashyap *et al.*

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.